

A Comment on the Strong Interactions of Color-Neutral Technibaryons

R. Sekhar Chivukula^{a,1}, Andrew G. Cohen^{a,2}, Michael Luke^{b,3} and Martin J. Savage^{b,4}

ABSTRACT

We estimate the cross section for the scattering of a slow, color-neutral technibaryon made of colored constituents with nuclei. We find a cross section of order $A^2 10^{-45} \text{ cm}^2$, where A is the atomic number of the nucleus. Even if technibaryons constitute the dark matter in the galactic halo, this is too small to be detected in future underground detectors.

10/92

^aBoston University, Department of Physics, 590 Commonwealth Avenue, Boston, MA 02215

^bDepartment of Physics 0319, University of California, San Diego, 9500 Gilman Drive, La Jolla, CA 92093-0319

¹sekhar@weyl.bu.edu

²cohen@andy.bu.edu

³luke@yukawa.ucsd.edu

⁴savage@thepub.ucsd.edu

If technicolor [1] is responsible for electroweak symmetry breaking, then technibaryons are an attractive “cold dark matter” candidate [2] [3] since their numbers relative to ordinary baryons may be *dynamically* determined [4] [5]. Current searches [6] for dark matter candidates using underground double- β decay detectors are able to exclude such technibaryons (with a mass of order 1 TeV) if they carry a weak charge and are also present with a density sufficient to explain the galactic halo. This exclusion is the result of coherent weak interactions in the scattering of the hypothetical technibaryon with nuclei.

It is nevertheless possible [2] that the lightest technibaryon has weak-isospin zero in which case the coherent weak interactions which produce a large scattering cross section with ordinary matter will be absent. For example, in the one-family model [7] with $N_{TC} = 4$ the lightest technibaryon may have a techniquark composition of either $UUDE$ or $UDDN$, where $\begin{pmatrix} U \\ D \end{pmatrix}$ and $\begin{pmatrix} N \\ E \end{pmatrix}$ are techniquark and technilepton doublets respectively. In this case [2], the lightest technibaryon has spin and isospin zero and is both color and electrically neutral; such a particle has no coherent weak interactions and would therefore be very hard to detect.

Nussinov [8] has noted that if the neutral technibaryon has colored constituents, then at distances less than approximately one Fermi (where one may ignore issues of confinement) it will have a nonzero chromoelectric polarizability. Using a non-relativistic quark model, Nussinov estimated the cross section for the scattering of a neutral technibaryon from the chromoelectric fields in a nucleon to be of order $A^2 10^{-38} \text{ cm}^2$, where A is the atomic number of the nucleus. While this cross section is too small to be observed currently, it is hoped that it may be large enough to be observed in future detectors.

In this note, we re-calculate the cross section for the scattering of a color-neutral technibaryon from nuclei without recourse to the quark model. For definiteness, we consider the spin-0 technibaryon ϕ of the one-family model discussed above.

Consider constructing an effective Lagrangian to describe the interaction of ϕ with the color field. The leading interaction comes from the two dimension seven operators

$$\frac{g^2}{(4\pi f)^3} (c_1 \phi_v^* \phi_v G^{a\mu\nu} G_{\mu\nu}^a + c_2 \phi_v^* \phi_v G^{a\mu\alpha} G_\alpha^{a\nu} v_\mu v_\nu) , \quad (1)$$

where $G_{\mu\nu}^a$ is the color field strength tensor, g is the color coupling constant, and f is the analog of f_π for the technicolor theory (which is approximately 125 GeV in the one-family model). Since the technibaryon is much heavier than Λ_{QCD} , strong interactions do not

change its velocity. It is then convenient to describe technibaryons with different velocities by distinct fields [9]. ϕ_v is related to the usual field ϕ by the field redefinition

$$\phi_v(x) = \sqrt{2m_\phi} e^{im_\phi v_\alpha x^\alpha} \phi(x), \quad (2)$$

where v^μ is the four-velocity of the technibaryon. Because of the phase redefinition (2), derivatives acting on ϕ_v bring down factors of the small “residual” momentum $k^\mu \equiv p^\mu - m_\phi v^\mu$. This prevents the derivative expansion from containing potentially troublesome terms proportional to $m_\phi v^\mu / 4\pi f \sim \mathcal{O}(1)$.

Classically, the interaction (1) corresponds to the interaction of induced chromoelectric and chromomagnetic dipole moments with the color field. Here, c_1 and c_2 are unknown strong interaction constants which, according to the rules of dimensional analysis [10], are expected to be of order 1. The operators and the coupling g are all understood to be renormalized at a scale of order 1 TeV. The matrix element of the second operator in (1) in nuclei may be related to the structure function conventionally called F_1 ; however it is suppressed relative to the first operator in (1) by a factor of $\sim \alpha_s(1 \text{ TeV})$, and we will consequently neglect it.

Up to small corrections of order $\alpha_s^2(1 \text{ TeV})$, we may rewrite the first term in (1) as

$$-c_1 \frac{\beta(g)}{4\pi f^3 b g} \phi_v^* \phi_v G^{a\mu\nu} G_{\mu\nu}^a, \quad (3)$$

where

$$\beta(g) \approx \frac{-bg^3}{16\pi^2} \quad (4)$$

is the beta function for the color coupling g . This reformulation is useful since the operator $\frac{\beta}{g} G^2$ is a renormalization group invariant and consequently its matrix elements between physical states are renormalization scale independent.

To calculate the cross section for scattering from a nucleon, we must evaluate the matrix element

$$\langle k, s | \frac{\beta}{g} G^2(q) | k', s \rangle \equiv -4m^2 S(q^2), \quad (5)$$

where $|k, s\rangle$ is a nucleon state (either p or n) of momentum k and spin s , $q = k - k'$ is the momentum transfer, m is the nucleon mass and $S(q^2)$ is a form factor. As noted by Voloshin and Zakharov [11], up to corrections suppressed by powers of the light quark masses, the operator $\frac{\beta}{g} G^2(0)$ is related to the generator of scale transformations; as noted

in [12] this amounts to differentiating with respect to the logarithm of Λ_{QCD} . The result is that, at zero momentum transfer

$$\langle k, s | \frac{1}{2} \frac{\beta}{g} G^2(0) | k', s \rangle = -2m^2 . \quad (6)$$

This implies $S(0) = 1$.

For collisions of technibaryons in the galactic halo (which have a velocity $v_{\text{halo}} \approx 10^{-3}c$) with nuclei in underground detectors, the momentum transfer is of order a few tens of MeV/c, while the scale of momentum over which S varies is of order 100 MeV/c [13]. Therefore, to a good approximation, we may replace $S(q^2)$ with 1. If the momentum transfer increases to a few hundred MeV/c this form factor will further suppress the cross section for elastic scattering.

We may now calculate the cross section for the scattering of a slow technibaryon from a nucleus in an underground detector

$$\sigma \simeq \frac{c_1^2 A^2 m^4}{(4\pi)^3 (1 + Am/M)^2 b^2 f^6} , \quad (7)$$

where M is the mass of the technibaryon and A is the atomic number of the nucleus. Here we have assumed that the technibaryon scatters coherently from all of the nucleons in a nucleus. Taking $m = 1$ GeV, and $b = 7$ (appropriate for six light quarks) we find a cross section $\sigma \approx A^2 10^{-45} \text{ cm}^2$, independent of the technibaryon mass for $Am/M \ll 1$. Unfortunately, this cross section is too small to be observed in any foreseeable future detector.

Our calculation gives a result approximately seven orders of magnitude smaller than the estimate given in [8]. This discrepancy can be explained by the fact that the large quark-model result is due to contributions from regions where the technibaryon is a distance of order $(1 \text{ TeV})^{-1}$ from a quark. However, this is a distance scale much smaller than the Compton wavelength of a “constituent” quark and the non-relativistic quark model is inappropriate. By contrast, since we are able to relate the cross section to the matrix element of $\frac{\beta}{g} G^2$, our calculation should be trustworthy.

Two issues remain. Firstly, we have so far only considered the coupling of our neutral technibaryon to gluons, without direct couplings to the light quarks. Such couplings will appear in our effective theory, but they will be suppressed either by powers of the extended technicolor scale [14] or by powers of $\alpha_s(1 \text{ TeV})$. For an ETC scale bigger than a few times 10 TeV, these contributions are negligible compared to the gluonic piece we have calculated.

Secondly, in other models of technicolor the lightest technibaryon may be a fermion rather than a boson. This makes no difference since the non-relativistic scattering amplitude will not distinguish a fermion from a boson.

Acknowledgements

We would like to thank Shmuel Nussinov for useful conversations and comments. R.S.C. and M.J.S. thank Robert Jaffe and Emil Mottola for organizing the *Sante Fe Workshop on Hadrons and Physics Beyond the Standard Model* where some of this work was completed. R.S.C. acknowledges the support of an Alfred P. Sloan Foundation Fellowship, an NSF Presidential Young Investigator Award, a DOE Outstanding Junior Investigator Award, and a Superconducting Super Collider National Fellowship from the Texas National Research Laboratory Commission. A.C acknowledges the support of a DOE Outstanding Junior Investigator Award. M.J.S. acknowledges the support of a Superconducting Super-collider National Fellowship from the Texas National Research Laboratory Commission.

This work was supported in part under NSF contracts PHY-9057173 and PHY-8958081 and DOE contracts DE-AC02-89ER40509, DE-FG02-91ER40676 and DE-FG03-90ER40546, and by funds from the Texas National Research Laboratory Commission under grants RGFY91B6 and FCF9219.

References

- [1] S. Weinberg, Phys. Rev. D **19** (1979) 1277;
L. Susskind, Phys. Rev. D **20** (1979) 2619.
- [2] S. Nussinov, Phys. Lett. **B165** (1985) 55.
- [3] R. S. Chivukula and T. Walker, Nucl. Phys. **B329** (1990) 445.
- [4] S. Barr, R. S. Chivukula, and E. Farhi, Phys. Lett. **B241** (1990) 387.
- [5] D. B. Kaplan, Phys. Rev. Lett. **68** (1992) 741.
- [6] D. O. Caldwell, *et al*, Phys. Rev. Lett. **61** (1988) 551 ;
S. P. Ahlen, *et al*, Phys. Lett. **B195** (1987) 603.
- [7] E. Farhi and L. Susskind, Phys. Rev. D **20** (1979) 3404.
- [8] S. Nussinov, Phys. Lett. **B279** (1992) 111.
- [9] H. Georgi, Phys. Lett. **240** (1990) 447.
- [10] H. Georgi and A. V. Manohar, Nucl. Phys. **B234** (1984) 189.
- [11] M. Voloshin and V. Zakharov, Phys. Rev. Lett. **45** (1980) 688.
- [12] R.S. Chivukula, A.G. Cohen, H.M. Georgi and A.V. Manohar Phys. Lett. **B222** (1989) 258.
- [13] see, for example, J. Engel, S. Pittel and P. Vogel, Bartol Preprint BA-92-23.
- [14] E. Eichten and K. Lane, Phys. Lett. **90B** (1980) 125;
S. Dimopoulos and L. Susskind, Nucl. Phys. **B155** (1979) 237.